

Comment on “New Methods for Old Coulomb Few-Body Problems”

V. S. Zotev

Los Alamos National Laboratory, Biophysics Group, MS D454, Los Alamos, NM 87545, USA

In this Comment on the above mentioned paper by F. E. Harris, A. M. Frolov, and V. H. Smith, we briefly review our contributions to development of new methods for solution of the Coulomb four-body problem. We show that our research group, headed by Prof. T. K. Rebane, had a priority in using the fully correlated exponential basis for variational calculations of four-body systems. We also draw attention to the fact that our group subsequently implemented a more advanced method, which uses highly efficient exponential-trigonometric basis functions for solution of the same problem.

The paper “New Methods for Old Coulomb Few-Body Problems” [1] deals with analytical and methodological issues involved in the implementation of two methods – the Hylleraas method and what the authors call the “exponential ansatz” approach. It is stated in the abstract that the second method “has only recently been implemented for four-body systems” [1]. The last section of the paper is entitled “Preliminary Results” [1].

The main purpose of this Comment is to call attention to our works in this field and show that the method using a large fully correlated exponential basis for solution of the Coulomb four-body problem was in fact implemented and successfully applied by our group ten years ago.

A general method for analytic evaluation of four-particle integrals was developed by Fromm and Hill in their well-known pioneering work [2] in 1987. This method makes it possible to perform direct variational calculations of four-body systems using exponential basis functions of all six interparticle separations. The first such calculation of two model systems was carried out by Rebane and Yusupov in 1992 [3]. Rebane showed in 1993 how the number of different types of four-body integrals that need to be computed can be reduced from 43 to only seven [4]. The first variational calculation of the positronium molecule $e^+e^-e^+e^-$ in the fully correlated exponential basis was performed by Rebane, Zotev, and Yusupov in 1996 [5]. It employed 90 symmetrized exponential functions. This work was followed by calculations of eight symmetric mesomolecules in the basis of 240 exponentials [6]. In 1998, the original method by Fromm and Hill was generalized by Zotev and Rebane to allow evaluation of four-particle integrals with complex exponential parameters. This enabled us to perform the first variational calculation of the positronium molecule and three other systems using the basis of extremely flexible exponential-trigonometric functions [7]. It was shown that a single exponential-trigonometric function is as efficient as seven exponentials in the case of $e^+e^-e^+e^-$, and is flexible enough to allow direct variational computation of even adiabatic four-particle systems such as the hydrogen molecule [7]. The exponential-trigonometric basis was initially introduced by our group in the three-body problem [8]. A detailed description of our methods using the exponential and exponential-trigonometric basis functions for direct variational solution of the Coulomb four-body problem was published in 2002 [9].

Harris, Frolov, and Smith implemented the exponential basis and reported a calculation of the positronium molecule with *two* exponential functions in 2003 [10]. They used a modified Fromm and Hill formula [11], and argued that the advantage of their approach was “better formal treatment of termwise singularities” [10]. They wrote in the discussed paper that their refinements enabled them to find “a two-configuration wavefunction with energy -0.513237 hartree, comparable to the result obtained by Rebane *et al.* using four configurations” [1].

Our extensive practical experience shows that the problem of termwise singularities is not serious, because parameters of basis functions can easily be chosen to avoid singularities in all the integrals to be computed. Our calculations of the positronium molecule [5] and mesomolecules [6] with the proper parameter selection exhibited perfect numeric stability and steady convergence.

The authors’ comparison of their two-function calculation and our four-function result is performed incorrectly. In their calculation [10], the two functions were optimized *simultaneously*. In our work [5], the basis was expanded gradually, and basis functions were optimized *one at a time*. Naturally, the simultaneous optimization provides better results. Moreover, our 1996 work included not just four, but a total of 90 exponential functions, and yielded a variational energy value of -0.515920 hartree [5].

Finally, we have to mention the following. Frolov and Smith [12], Frolov [13], Frolov and Bailey [14], when discussing the Fromm and Hill four-body integrals in their 2001-03 articles, did not reference our works at all. Similarly, Frolov made no reference to our original works [8] in his 1998-03 papers, which used the exponential-trigonometric basis in the three-body problem [15]. In the discussed paper [1], Harris, Frolov and Smith cite our 1996 calculation of the positronium molecule [5], but do not reference any subsequent publications [6, 7, 9], particularly the 2002 article [9], which provides a detailed review of our methods. Frolov and Smith follow the same line in another recent paper [16]. The most advanced and accurate method, which uses highly flexible exponential-trigonometric functions for calculations of four-particle systems with arbitrary particle masses [7, 9], has never been mentioned by Harris, Frolov, and Smith. Even the latest and most comprehensive review article by Harris [17] makes no explicit mention of this method. We leave these facts without comment.

-
- [1] F. E. Harris, A. M. Frolov, and V. H. Smith, Int. J. Quantum Chem. **100**, 1086 (2004). The authors refer to this article as a “review of methods applicable to few-body problems” (*ibid.* p.1090).
- [2] D. M. Fromm and R. N. Hill, Phys. Rev. A **36**, 1013 (1987).
- [3] T. K. Rebane and O. N. Yusupov, Opt. Spektrosk. **73**, 875 (1992) [Opt. Spectrosc. **73**, 523 (1992)]. Values of four-body integrals for this work were provided by the Fromm and Hill reserach group.
- [4] T. K. Rebane, Opt. Spektrosk. **75**, 945 (1993) [Opt. Spectrosc. **75**, 557 (1993)]. See also T. K. Rebane, Opt. Spektrosk. **93**, 908 (2002) [Opt. Spectrosc. **93**, 837 (2002)].
- [5] T. K. Rebane, V. S. Zotev, and O. N. Yusupov, Zh. Eksp. Teor. Fiz. **110**, 55 (1996) [JETP **83**, 28 (1996)].
- [6] V. S. Zotev and T. K. Rebane, Opt. Spektrosk. **85**, 935 (1998) [Opt. Spectrosc. **85**, 856 (1998)].
- [7] V. S. Zotev and T. K. Rebane, Yad. Fiz. **63**, 46 (2000) [Phys. At. Nucl. **63**, 40 (2000)]. This paper was accepted in 1998, but published, for technical reasons, in 2000.
- [8] T. K. Rebane and O. N. Yusupov, Zh. Eksp. Teor. Fiz. **98**, 1870 (1990) [JETP **71**, 1050 (1990)]; V. S. Zotev and T. K. Rebane, Opt. Spektrosk. **77**, 733 (1994) [Opt. Spectrosc. **77**, 654 (1994)].
- [9] V. S. Zotev and T. K. Rebane, Phys. Rev. A **65**, 062501 (2002).
- [10] F. E. Harris, A. M. Frolov, and V. H. Smith, J. Chem. Phys. **119**, 8833 (2003).
- [11] F. E. Harris, Phys. Rev. A **55**, 1820 (1997).
- [12] A. M. Frolov and V. H. Smith, J. Chem. Phys. **115**, 1187 (2001).
- [13] A. M. Frolov, Phys. Rev. A **64**, 042503 (2001).
- [14] A. M. Frolov and D. H. Bailey, J. Phys. B **36**, 1857 (2003).
- [15] A. M. Frolov, Phys. Rev. A **57**, 2436 (1998); Phys. Rev. E **62**, 8740 (2000); Phys. Rev. E **64**, 036704 (2001); Phys. Rev. E **65**, 046705 (2002); Phys. Rev. A **67**, 064501 (2003).
- [16] A. M. Frolov and V. H. Smith, J. Phys. B **37**, 2917 (2004). The authors state in this paper that the methods based on the Fromm and Hill formula “are not, probably, highly accurate” (*ibid.* p.2930).
- [17] F. E. Harris, Advances in Quantum Chemistry **47**, 129 (2004).